

**HURLEY DISEASE - SHORT COMMENT**Anca Chiriac<sup>1</sup>, Liliana Foia<sup>2</sup>, Tudor Pinteala<sup>3</sup>, Anca E. Chiriac<sup>4</sup><sup>1</sup>*CMI Dermatology-Iasi,Romania*<sup>2</sup>*Univ.of Medicine Gr T Popa, Biochemistry Department, Iasi-Romania*<sup>3</sup>*Imperial College of London, UK*<sup>4</sup>*Univ.of Medicine Gr T Popa Iasi-Romania***Source of Support:**  
Nil**Competing Interests:**  
None declared**Corresponding author:** Anca Chiriac, MD PhD[ancachiriac@yahoo.com](mailto:ancachiriac@yahoo.com)

Our Dermatol Online. 2012; 3(4): 363

Date of submission: 06.06.2012 / acceptance: 13.06.2012

**Cite this article:***Anca Chiriac, Liliana Foia, Tudor Pinteala, Anca E. Chiriac: Hurley disease. Our Dermatol Online. 2012; 3(4): 363***Short comment to a previous article:***Chiriac A, Foia L, Pinteala T, Chiriac AE: Acne inversa (Hurley clinical stage ii): case report. Our Dermatol Online. 2011; 2(4): 215-216.*

Sir,

The patient refused the surgery and we recommended him resorcinol 30% in vaseline, vitamin C high doses orally (2g/day) and a two-month treatment with Sultametoxazol 4cp/day (Biseptol, Cotrimoxazol) plus Rifampicine 300 mg/day, with very good results (as seen in the picture).

Unfortunately we were obliged to stop the systemic medication due to the gastro-intestinal side effects, although the laboratory investigations were all within normal limits.

**REFERENCES**

1. Chiriac A, Foia L, Pinteala T, Chiriac AE: Acne inversa (Hurley clinical stage II): case report. Our Dermatol Online. 2011;2:215-6.



Figure 1.




Figure 2.